



**PubMed**

National  
Library  
of Medicine

PubMed

Nucleotide

Protein

Genome

Structure

PMC

Taxonomy

OMIM

Books

Search PubMed

for

☒ Limits

Preview/Index

History

Clipboard

Details

Go Clear

About Entrez

Text Version

Display

Abstract

Show: 20

Sort

Send to

Text

☐ 1: Nature 1994 Apr 28;368(6474):856-9

Related Articles: Links

Comment in:

- Nature. 1994 Apr 28;368(6474):812

### Antigen-specific human antibodies from mice comprising four distinct genetic modifications.

Lonberg N, Taylor LD, Harding FA, Trounstein M, Higgins KM, Schramm SR, Kuo CC, Mashayekh R, Wymore K, McCabe JG, et al.

GenPharm International, Mountain View, California 94043.

PubMed Services  
Journals Database  
MeSH Browser  
Single Citation Matcher  
Batch Citation Matcher  
Clinical Queries  
LinkOut  
Cubby

Related Resources  
Order Documents  
NLM Gateway  
TOXNET  
Consumer Health  
Clinical Alerts  
ClinicalTrials.gov  
PubMed Central  
Privacy Policy

Human sequence monoclonal antibodies, which in theory combine high specificity with low immunogenicity, represent a class of potential therapeutic agents. But nearly 20 years after Kohler and Milstein first developed methods for obtaining mouse antibodies, no comparable technology exists for reliably obtaining high-affinity human antibodies directed against selected targets. Thus, rodent antibodies, and in vitro modified derivatives of rodent antibodies, are still being used and tested in the clinic. The rodent system has certain clear advantages; mice are easy to immunize, are not tolerant to most human antigens, and their B cells form stable hybridoma cell lines. To exploit these advantages, we have developed transgenic mice that express human IgM, IgG and Ig kappa in the absence of mouse IgM or Ig kappa. We report here that these mice contain human sequence transgenes that undergo V(D)J joining, heavy-chain class switching, and somatic mutation to generate a repertoire of human sequence immunoglobulins. They are also homozygous for targeted mutations that disrupt V(D)J rearrangement at the endogenous heavy- and kappa light-chain loci. We have immunized the mice with human proteins and isolated hybridomas secreting human IgG kappa antigen-specific antibodies.

PMID: 8159246 [PubMed - indexed for MEDLINE]

Display

Abstract

Show: 20

Sort

Send to

Text